

THE ASSESSMENT OF THE SENSORY QUALITY OF THE SAUERKRAUT FROM ORGANIC AND CONVENTIONAL FARMING

Summary

Sauerkraut is a popular food product in Poland. The consumption of sauerkraut, according to different sources, ranges from 4 to 5 kg/person/year. The objective of this study was to provide information on the nutritional and health properties of sauerkraut and to assess some selected distinctive features of sensory quality of sauerkraut from organic and conventional farming. The hypothesis of the study assuming that organic sauerkraut has a better tastiness, has been verified positively. The assessed types of sauerkraut have significantly different taste and aroma, the consumers have valued the organic sauerkraut significantly higher than the sauerkraut from conventional farming. A high level of positive correlation (0.994) between the organic and conventional products has been determined.

Key word: sauerkraut, sensory evaluation, consumer, organic food

KONSUMENCKA OCENA JAKOŚCI SENSORYCZNEJ KAPUSTY KISZONEJ Z PRODUKCJI EKOLOGICZNEJ I KONWENCJONALNEJ

Streszczenie

Kapusta kiszona jest bardzo popularnym produktem żywnościowym w Polsce. Spożycie to kształtuje się według różnych źródeł od 4 do 5 kg/osobę/rok. Celem pracy było przybliżenie wiedzy na temat właściwości odżywczych i zdrowotnych kapusty kiszonej oraz ocena wybranych wyróżników jakości sensorycznej kapusty kiszonej otrzymanej z kapusty uprawianej w sposób ekologiczny i konwencjonalny. Postawiona w pracy hipoteza o wyższej smakowitości ekologicznej kapusty kiszonej została zweryfikowana pozytywnie. Oceniane kapusty kiszone znacznie różniły się smakiem i aromatem, zdecydowanie wyżej konsumenci ocenili kapustę z produkcji ekologicznej. Stwierdzono bardzo wysoką korelację dodatnią (0,994) pomiędzy produktami ekologicznymi a konwencjonalnymi.

Słowa kluczowe: kapusta kiszona, ocena sensoryczna, konsument, żywność ekologiczna

1. Introduction

In the recent years, the consumers have been paying more and more attention to the quality of food. It results from, among other factors, a wide range of food products on the market and the difficulty in choosing the right product. However, the growing awareness of the consumers regarding the nutritional quality and value of food has become a new factor. Increased demand for organic food, which is considered to be healthy, tasty and of high nutritional value, has been observed. The production of organic food based on special norms, is regarded as environment-friendly.

The cruciferous vegetables (*Brassicaceae*) belong to the plant products which are the basic food products for the citizens of many European and Asian countries. The vegetables of this family are produced in various agricultural systems and with the use of various organic methods.

The cruciferous vegetable family (*Brassicaceae*) includes a great number of vegetables, such as cabbage, cauliflower or kale. There are numerous varieties of cabbage, but one of the greatest significance is the headed cabbage, including the following varieties: the white cabbage (*Brassica oleracea* L. var. *capitata* f. *alba*), red cabbage (*B. oleracea* var. L. *capitata* f. *rubra*) and Brussel sprout (*B. oleracea* L. var. *gemmifera* Zenk.). Cabbage is an essential element of a human diet, especially in Europe and Asian countries. The vegetables of *Brassicaceae* species have been gaining popularity as they contain numerous compounds which have a positive influence on the human health, including glucosinolates and polyphenols [8, 19]. The glucosinolates are ac-

companied by myrosinase, an enzyme responsible for the glucosinolates hydrolysis to numerous bio-active compounds such as thiocyanates and isothiocyanates, which inhibit the growth of microorganisms and have anticarcinogenic properties [11]. The most significant polyphenolic compounds are the derivatives of quercetin and kaempferol [24]. The cruciferous vegetables are also the source of vitamins C and E, β -Carotene, luteine and DL- α -tocopherol. It has been proved that the compounds are powerful antioxidants and they protect cells against oxidative stress. Therefore, they may prevent from numerous chronic diseases such as neoplasms, cardio-vascular diseases and diabetes [19].

The presence of bio-active compounds in the *Brassicaceae* family and their anti-oxidant properties depends on numerous factors, including the genetic traits (the species, variety, part of the plant), climate conditions, the soil composition and fertility, agronomic treatments, the harvesting conditions, storage and processing [1, 9, 22]. The common methods of the processing of *Brassicaceae* include blanching, boiling and freezing. One of the most commonly used methods of cabbage processing consists in fermentation. Sauerkraut is a very popular and often consumed vegetable in Poland, its consumption is about 4 kg/year/person [25]. The cabbage both from organic and conventional farming is used for the production of sauerkraut.

Generally it shall be deemed that organic food is more healthy and tasty as no synthetic fertilisers, chemical products as pesticides and herbicides, are used in the production process, whereas the cabbage processing involves natural additives [26]. The processing itself is subject to develop-

ment and the organic sauerkraut has been present on the market for several years.

The objective of the study was to assess selected distinctive features of sensory quality of sauerkraut from organic and conventional farming.

The hypothesis has been presented in this study that organic sauerkraut has a higher level of tastiness than sauerkraut from the conventional farms.

2. Cabbage fermentation technology

The technology of cabbage fermentation is very simple. The ingredients used in the fermentation process are: cabbage, common salt and water. The certified cabbage from organic farms is used in the production of organic sauerkraut and the fermentation process runs spontaneously.

A fresh headed cabbage *Brassicaoleracea* L. var. *capitata* is the best variety of cabbage to be used for fermentation. A stored cabbage loses its bio-active compounds depending on the storage conditions and period. The studies have confirmed that even the storage of cabbage in controlled atmosphere causes the amount of vitamin C and chlorophyll to decrease [10, 14, 30]. The research has shown that the cabbage from organic farming contained more bio-active compounds than the cabbage from conventional farming, but the storage stability of those compounds in different farming systems was diversified [6].

The fermentation process involves the bacteria present on the surface of crushed cabbage leaves and bacteria present in the air of the room where the process takes place. The fermentation is started by the heterofermentative bacteria of lactic acid, such as *Leuconostoc mesenteroides* which produces mainly acetic acid and lactic acid. Further fermentation takes place with the use of *Lactobacillus plantarum* [3, 31].

The spontaneous fermentation might cause differences in the composition of microbiota, which may be the reason of the quality change of sauerkraut. For the standardization process the starter cultures are used in the industrial production [29]. Most frequently they include the bacteria cultures isolated from sauerkraut, such as *Leuconostoc mesenteroides*, *Lactobacillus plantarum*, *Lactobacillus casei* and *Lactococcus lactis*. The studies with these starter cultures have been conducted in China, among other countries. The bacteria used in the industrial fermentation were isolated from sauerkraut which underwent a traditional fermentation process. This sauerkraut should be differentiated from kimchi, the sauerkraut produced in some of the regions of China, but mainly in Korea and Taiwan [4, 23]. In the sauerkraut produced with the use of starter cultures, the fermentation process is controlled, and the quality of product is standardized. In the spontaneous fermentation the quality depends on many factors [29].

In order to improve the taste and the aroma, the additions of dill, carrots and apples are used in the fermentation process. These additives are more often used in the organic production of sauerkraut. Their role is to improve the taste of the food. A great number of spices also have a therapeutic effect and facilitates digestion. However, the microbiological purity of these spices is a great problem. The collection of spices and preparation technology are traditional and fosters the habitation of bacteria, mould and fungi in the dried plants mostly. The general amount of microorganisms in spices may come up to 10^9 in 1 g. In the industrial production of spices the radiation sterilization is used [4, 23]. An excessive addition of spices in the organic production of

sauerkraut may interrupt the spontaneous fermentation and result in a "wild" fermentation.

The direction of fermentation and the quality of the fermented cabbage are also influenced by the amount of salt. The research conducted in China has shown that the amount of lactic acid has increased with the increase of salt concentration. The higher concentrations of salt successfully inhibited the growth of fungi and *E. coli*. For comparison, a high concentration of salt slowed down the maturity of cabbage and inhibited the metabolism of the lactic acid bacteria [32].

The content of sodium in sauerkraut is essential as an excessive amount of sodium is associated with an increased blood pressure and it is a risk factor for cardio-vascular diseases. This is the reason for the calls of the health organizations and consumers organizations for reducing the amount of sodium in food. The actions concerning this problem are also directed towards the process of cabbage fermentation. The studies have shown that the amount of salt in sauerkraut may range from 15.0 to 22.5 g kg⁻¹ NaCl [23]. Numerous experiments resulting in the reduction of the amount of NaCl in the cabbage fermentation process have shown that the reduction of the amount NaCl to 5% and the addition of various chemical compounds such as KCl, HCl, MgSO₄, SiSO₂ result in the reduction of the salt amount, however it changes the taste of the sauerkraut, which must be concealed by spices. Moreover, the cabbage becomes less crunchy [31].

Another direction of research on cabbage fermentation included actions aiming to enrich this product with selenium compounds. The role of selenium is to prevent from chronic diseases related to oxidative stress. According to Penas et al [18], the addition of selenium salt has increased the content of glucosinolates and se-methylselenocysteine improving the antioxidative ability and potential anti-inflammatory properties of sauerkraut.

3. Health properties and the presence of anti-nutrients in sauerkraut

The health properties of sauerkraut are connected first of all with the content of such compounds like polyphenol compounds, vitamins C, E, K, carotenoids, flavonoids, sulphides, fiber, selenium, zinc, manganese, potassium, iron and the others. The compounds characterized with high antioxidant and antiradical activity play particular role among these compounds.

The sauerkraut processing procedures in an increased temperature such as blanching, pasteurization, sterilization cause the antioxidant properties of sauerkraut to decrease. On the other hand, the fermentation of sauerkraut causes increase in the content of antioxidant compounds and vitamin C [5, 17].

Glucosinolates draw the biggest attention out of all the substances occurring in cruciferous vegetables. Until not long ago these compounds had been considered as anti-nutrients [13, 28]. They have a very complex chemical structure, over 200 compounds have been discovered until now, and they hydrolyze to isothiocyanates, thiocyanate, nitriles and indoles. These compounds have a very unpleasant taste and smell, they cause irritation of the alimentary tract, they also have goitrogenic properties. They get decomposed during cooking and they evaporate with steam. The latest research has confirmed that glucosinolates have the ability to inhibit

cancer processes. Fresh and sour red cabbage and fresh white cabbage and sauerkraut indicate the highest antioxidant activity out of cruciferous vegetables [12]. During fermentation the decomposition of glucosinolates takes place and the intensity of this process depends on the fermentation temperature and the population of lactic acid bacteria [15].

Besides the anticancer effect, sauerkraut has a positive influence on blood pressure, immunity system, it enhances metabolism due to the content of fiber, it has weight loss properties, reduces the level of cholesterol and has effects on lipid metabolism, it increases iron absorption [25].

Following to the microorganisms' functioning in food, free amines having strong effects on the organism, called biogenic amines, come into being. This process also occurs during sauerkraut fermentation. It has been found out that histamine is created in sauerkraut in the biggest amount. The content of histamine in fresh vegetables depends on the type, species, degree of maturity and on processing and storing of the products. Histamine indicates psychoactive and vasoactive properties which can induce toxicological effects.

Histamine in high concentration causes headache. Other biogenic amines include: tyramine inducing migraines and arterial hypertension, spermine and spermidine which are less toxic, however they enhance the negative effect of histamine. Studies have confirmed that the microorganisms used in the fermentation process have effect on the amount and type of the created amines. The total concentration of the biogenic amines in sauerkrauts was lower when the fermentation was conducted with *L. mesenteroides* rather than with *L. plantarum* [16].

4. Material and methods

The sauerkraut from organic and conventional farming was the material for studies. In both cases the cabbage was subject to fermentation in special fermentation tanks, next it was put to jars in order to undergo pasteurization. The organic product was properly marked with a control unit certificate. The conventional product had the basic information regarding the product and the supplier, however, no information was given whether any other additions had been used in the fermentation process, such as acid addition, which is a common procedure in the conventional farming and it provides microbiological protection against the growth of any undesired microorganisms.


The samples of sauerkraut from the organic and conventional farming were coded. The packaging compliance was assessed in terms of the requirements of the Regulation of the Ministry of Agriculture and Rural Development on Labelling of Foodstuffs for Particular Nutritional Purposes of 23rd December 2014 and of The Act of 25 August 2006 on Food and Nutrition Safety [20, 21, 27]. The labels of the examined products included the basic information such as: name of the product, net weight, name of the manufacturer, batch number, ingredients and minimum expiry date, whereas the organic sauerkraut also had the relevant organic food certification marks. All captions were in Polish, they were legible and permanent. There was no information about the presence of allergens on the labels of either the organic or conventional product.

A sensory analysis was carried out by means of a five point grading scale for quality factors. The following parameters were assessed: the appearance, colour, aroma,

taste [2]. The scaling method which allowed for a digital expression of the examined parameters (taste, aroma, appearance, colour) was used for examining the quality of the sauerkraut, taking into account the ranking scale of particular factors. The organoleptic analysis was performed according to Five Point Scale instituted by Tilgner (1 point means disqualification whereas 5 means very good). MS Excel sheet and Statistica 12 package for data statistical analysis were used for the calculation of the final score of the organoleptic analysis. The following significance coefficients were assumed: 0,5 – taste; 0,3 – aroma; 0,1 – appearance; 0,1 – colour [7].

5. Test results

A consumer-like assessment of sauerkrauts has been performed and participated by 58 young women aged 20-26. The persons assessing the sauerkraut did not eat or drink directly before tasting, nor did they smoke cigarettes, which could have affected their objective sensory perception. The samples, all formerly cooled down to 6°C, were served on white plates directly before the assessment. Following to the assessment the average score from 1.25 to 5 points was awarded for the organically farmed sauerkraut whereas the score from 1.25 to 4.5 was awarded to the conventionally farmed sauerkraut (Fig. 1).



Source: own work / Źródło: opracowanie własne

Fig. 1. Organoleptic assessment of sauerkraut from organic and conventional farming

Rys. 1. Ocena organoleptyczna kapusty kiszonej z gospodarki ekologicznej i konwencjonalnej

The results were subjected to statistical analysis, by means of one factor variance analysis, at $p=0.05$ significance level, using the data statistical analysis package Statistica 12. No statistically significant difference ($p<0.05$) was indicated in the assessment of sauerkraut from conventional and organic farming ($p=0.094$).

It is difficult to discuss the results obtained from the study due to the fact that no research in terms of the sensory assessment of sauerkraut from organic and conventional farming has been conducted so far.

The assessed sauerkrauts significantly differed from each other in terms of taste and aroma, the consumers valued the sauerkraut from organic farming remarkably higher. A very high positive correlation (0.994) has been confirmed


between the organic and conventional products (Tab. 1, Fig. 2).

Table 1. The sensory quality assessment of sauerkraut from organic and conventional farming

Tab. 1. Ocena jakości sensorycznej kapusty kiszzonej z gospodarki ekologicznej i konwencjonalnej

Subject of the evaluation	Organic [$\bar{X} \pm SD$]	Conventional [$\bar{X} \pm SD$]	Grade
Appearance	3.76 ± 1.16	3.67 ± 1.15	0.1
Colour	3.52 ± 1.16	3.69 ± 0.90	0.1
Aroma	3.60 ± 1.09	3.19 ± 1.08	0.3
Taste	3.60 ± 1.23	3.24 ± 1.30	0.5
Overall rating	3.61 ± 0.97	3.31 ± 0.92	

Source: own work / Źródło: opracowanie własne



Source: own work / Źródło: opracowanie własne

Fig. 2. Quality factors of the organoleptic assessment of sauerkraut from organic and conventional farming

Rys. 2. Czynniki jakościowe oceny organoleptycznej kapusty kiszzonej z gospodarki organicznej i konwencjonalnej

6. Summary

Sauerkraut is a valued food product in Poland and in many countries of Europe and Asia. This thesis presents the actual status of knowledge about the dietary and nutritional importance of sauerkraut. The difference between the organic and conventional sauerkraut lies in the method of its fermentation. The fermentation is spontaneous in the organic process whereas it is controlled by starter cultures in the conventional process.

The approach to anti-nutrients contained in cruciferous vegetables has changed following to the research carried out by many scientists. Sauerkraut is characterized by the higher content of vitamin C and glucosinolates than fresh cabbage. In the course of myrosinase glucosinolates get decomposed to isothiocyanates, thiocyanates, nitriles and indoles. The fermentation process increases the content of these compounds. It has been indicated that the products of glucosinolates decomposition, together with polyphenols, determine the antioxidant properties of sauerkrauts.

Presently, the research on sauerkraut is oriented on increasing the sauerkraut value by reducing the use of salt during the fermentation process and using the addition of

other substances, such as selenium, which might cause that sauerkraut would be considered as functional food.

Polish consumers highly rate sauerkraut and they use it for fresh meals, sauerkraut soups and for the traditional dish called *bigos*.

Sauerkraut obtained from the cabbage grown on organic and conventional farms was subject to assessment in this study. Young women who consumed sauerkraut very frequently and willingly valued the sauerkraut from organic farming higher, pointing at its better taste and aroma. They also paid attention whether the texture that is crispiness of the organic sauerkraut was higher. The price was a factor limiting the frequent consumption of sauerkraut of organic origin.

7. References

- [1] Aires A., Carvalho R., Rosa E.: Glucosinolate composition of Brassica is affected by Postharvest, Food processing and myrosinase activity. *J. Food Process. Preserv.*, 2012, 36, 214-224.
- [2] Babicz-Zielińska E., Rybowska A., Obniska W.: *Sensoryczna ocena jakości żywności*, Akademia Morska w Gdyni, Gdynia 2008, 56-58.

- [3] Beganović J., Kos B., Pavunc A.B., Uroić K., Jokić M., Šušković J.: Traditionally produced sauerkraut as source of autochthonous functional starter cultures. *Microbiol. Res.*, 2014, 169, 623-632.
- [4] Benerjee M., Sarkar K.: Microbiological quality of some retail spices in India. *Food Research International*, 2003; 36: 469-474.
- [5] Ciska, E., Honke, H.: Effect of the pasteurization process on the content of ascorbigen, indole-3-carbinol, indole-3-acetonitrile, and 3,30-diindolylmethane in fermented cabbage. *J. Agric. Food Chem.*, 2012, 60, 3645-3649.
- [6] Gałązka-Czarnecka I., Krala L.: Stabilność przechowalnicza zawartości związków w wybranych owocach i warzywach z upraw ekologicznych i konwencjonalnych. *Chłodnictwo*, 2012, t. XLVII, 7-8, 50-54.
- [7] Gawęcka J., Jędryka T.: Analiza sensoryczna. Wybrane metody i przykłady zastosowań. Akademia Ekonomiczna w Poznaniu, 2001, 57.
- [8] Halkier B.A., Gershenzon J.: Biology and biochemistry of glucosinolates. *Annu. Rev. Plant Biol.*, 2006, 57, 303-333.
- [9] Korus A., Lisiewska Z.: Effect of preliminary processing and method of preservation on the content of selected antioxidative compounds in kale (*Brassica oleracea* L. var. *pacephala*) leaves. *Food Chem.*, 2011, 129, 149-154.
- [10] Krala L., Witkowska M.: Kinetyka degradacji witaminy C w warzywach kapustnych przechowywanych w kontrolowanej atmosferze. *Przem. Ferment. Owoc-Warzyw.*, 2005, 12, 30-32.
- [11] Kuroiwa, Y., Nishikawa, A., Kitamura, Y., Kanki, K., Ishii, Y., Umemura, T., Hirose, M.: Protective effects of benzyl iso-thiocyanate and sulforaphane but not resveratrol against initiation of pancreatic carcinogenesis in hamsters. *Cancer Lett.*, 2006, 241, 275-280.
- [12] Kusznierevicz B., Piasek A., Lewandowska J., Bartoszek A.: Właściwości przeciwnowotworowe kapusty białej. *Zywność Nauka Technologia Jakość*, 2007, 6 (55), 20-34.
- [13] Kwiatkowska E., Bawa S.: Znaczenie substancji uznanych za antyodżywcze w profilaktyce chorób cywilizacyjnych. *Med. Rodzinna*, 2007, 2, 36-40.
- [14] Lee S.K., Kedar A.A.: Preharvest and post harvest factors influencing vitamin C content of horticultural crops. *Postharvest Biology and Biotechnology*, 2000, 20 (2), 207-220.
- [15] Palani K., Harbaum-Piayd B., Meske D., Kepple J.K., Bockelmann W., Heller K.J., Schwarz K.: Influence of fermentation on glucosinolates and glucobrassicin degradation products in sauerkraut. *Food Chem.*, 2016, 190, 755-762.
- [16] Peñas E., Frias J., Sidro B., Vidal-Valverde C.: Impact of fermentation conditions and refrigerated storage on microbial quality and biogenic amine content of sauerkraut. *Food Chem.*, 2010, 123, 143-150.
- [17] Peñas E., Limón R.I., Vidal-Valverde C., Frias J.: Effect of storage on the content of indole-glucosinolate breakdown products and vitamin C of sauerkrauts treated by high hydrostatic pressure. *LWT - Food Sci. Technol.*, 2013, 53, 285-289.
- [18] Peñas E., Martínez-Villaluenga C., Frias J., Sánchez-Martínez M.J., Pérez-Corona M.T., Madrid Y., Cámara C., Vidal-Valverde C.: Se improves indole glucosinolate hydrolysis products content, Se-methylselenocysteine content, antioxidant capacity and potential anti-inflammatory properties of sauerkraut. *Food Chemistry*, 2012, 132, 907-914.
- [19] Podśędek A.: Natural antioxidants and antioxidant capacity of Brassica vegetables: A review. *LWT- Food Sci. Technol.*, 2007, 40, 1-11.
- [20] Regulation (eu) no 1169/2011 of the European Parliament and of the council of 25 October 2011 on the provision of food information to consumers, amending Regulations (EC) No 1924/2006 and (EC) No 1925/2006 of the European Parliament and of the Council, and repealing Commission Directive 87/250/EEC, Council Directive 90/496/EEC, Commission Directive 1999/10/EC, Directive 2000/13/EC of the European Parliament and of the Council, Commission Directives 2002/67/EC and 2008/5/EC and Commission Regulation (EC) No 608/2004. <http://eur-lex.europa.eu/legalcontent/EN/TXT/PDF> [dostęp: 10.05.2017].
- [21] Rozporządzenie Ministra Rolnictwa i Rozwoju Wsi w sprawie znakowanie poszczególnych rodzajów środków spożywczych z dnia 23 grudnia 2014 roku. *Dz. U.* 08.01.2015 r., poz. 29.
- [22] Šamec, D., Piljac-Žegarac, J., Bogović, M., Habjanič, K., Grúz, J.: Antioxidant activity of white (*Brassica oleracea* L. var. *capitata*) and Chinese (*Brassica rapa* L. var. *pekinensis* (Lour.)) cabbage: The influence of development stage, cultivar choice and seed selection. *Sci. Hortic.*, 2011, 128, 78-83.
- [23] Schweiggert U., Carle R., Schieber A.: Conventional and alternative processes for spice production - a review. *Trends Food Sci. Technol.*, 2007, 18, 260-268.
- [24] Singh J., Upadhyay A.K., Prasad K., Bahadur A., Rai M.: Variability of carotens, vitamin C, E and phenolics in Brassica vegetables. *J. Food Comp. Anal.*, 2007, 20, 106-120.
- [25] Szwejda-Grzybowska J.: Właściwości prozdrowotne kiszzonej kapusty. *Przem. Ferment. Owoc-Warzyw.*, 2010, 5, 22-23.
- [26] Śmiechowska M.: Organic food in the opinion of Polish consumers on the background of European trends. *Polish Journal of Commodity Science*, 2007, 4(13), 44-55.
- [27] Ustawa z dnia 25 sierpnia 2006 r. o bezpieczeństwie żywności i żywienia. [Dz. U. 2006, Nr 171, poz. 1225]
- [28] Vig A.P., Rampal G., Thind T.S., Arora S.: Bio-protective effects of glucosinolates - A review. *LWT - Food Sci. Technol.*, 2009, 42, 1561-1572.
- [29] Wiander B., Ryhänen E.-L. Laboratory and large-scale fermentation of white cabbage into sauerkraut and sauerkraut juice by using starters in combination with mineral salt with a low NaCl content. *Eur. Food Res. Technol.*, 2005, 220, 191-195.
- [30] Witkowska M., Krala L.: Wpływ kontrolowanej atmosfery na jakość kapusty włoskiej przechowywanej chłodniczo. *Chłodnictwo*, 2006, t. XLI, 4, 52-56.
- [31] Wolkers-Rooijackers J.C.M., Thomas S.M., Nout, M.J.R.: Effects of sodium reduction scenarios on fermentation and quality of sauerkraut. *LWT - Food Sci. Technol.*, 2013, 54, 383-388.
- [32] Xiong T., Li J., Liang F., Wang Y., Guan Q., Xie M.: Effects of salt concentration on Chinese sauerkraut fermentation. *LWT - Food Sci. Technol.*, 2016, 69, 169-174.